

Oral presentation

Open Access

Tolerability and efficacy of the trifunctional antibody removab® (anti-EpCAM x anti-CD3) in patients with malignant ascites due to ovarian cancer: Results of a phase I/II study

P Wimberger*¹, A Burges², V Gorbounova³, H Sommer⁴, B Schmalfeldt⁵, J Pfisterer⁶, M Lichinitser³, A Makhson⁷, M Ströhlein⁸, W Eiermann⁹, M Biakhov¹⁰, V Moiseenko¹¹, A du Bois¹² and R Kimmig¹

Address: ¹Department of Obstetrics and Gynecology, University Hospital, Essen, Germany, ²Department of Obstetrics and Gynecology, University Hospital Munich/Grosshadern, Munich, Germany, ³Research Center for Oncology, Moscow, Russia, ⁴Department of Obstetrics and Gynecology, University Hospital Munich-City, Munich, Germany, ⁵Department of Gynecology, University Hospital Technical University, Munich, Germany, ⁶Department of Obstetrics and Gynecology, University Hospital, Kiel, Germany, ⁷Moscow Oncology Clinical Hospital, Moscow, Russia, ⁸Department of Surgery, University Hospital Munich/Grosshadern, Munich, Germany, ⁹Frauenklinik vom Roten Kreuz, Munich, Germany, ¹⁰Central Clinical Hospital of Ministry, Moscow, Russia, ¹¹St. Petersburg Scientific Oncology Institute, St. Petersburg, Russia and ¹²Department of Gynecology and gynecological Oncology, Dr. Horst-Schmidt-Kliniken, Wiesbaden, Germany

Email: P Wimberger* - pauline.wimberger@medizin.uni-essen.de

* Corresponding author

from Association for Immunotherapy of Cancer: Cancer Immunotherapy – 2nd Annual Meeting
Mainz, Germany, 6–7 May 2004

Published: 1 July 2004

Received: 28 April 2004

Cancer Cell International 2004, **4**(Suppl 1):S3

This article is available from: <http://www.cancerci.com/content/4/S1/S3>

Introduction

Malignant ascites in patients with gynecological malignancies is associated with poor prognosis and poor quality of life. The bispecific trifunctional antibody removab® (anti-EpCAM x anti-CD3) belongs to a new class of intact antibodies that has been developed for targeted therapy of epithelial tumors. The two binding sites of removab® are directed against epithelial tumor cells (EpCAM⁺) and T-cells (CD3⁺) thus recruiting T-cells in the direct environment of tumor cells. Simultaneously, mediated by the intact Fc-fragment removab® binds to FcγI/IIIR⁺ accessory cells (e.g. macrophages, natural killer cells, dendritic cells) that are mandatory for the induction of a tumor-specific immune response.

Patients and methods

In an open-label multicenter phase I/II-dose escalating study, a total of 23 patients with ovarian cancer and symptomatic ascites at FIGO stage III-IV were treated intraperitoneally with removab®. The patients had received a median of 3 (1–8) previous therapies. Their mean age was 62 (42–80) years. The treatment consisted of up to 5 intra-

peritoneal applications of the antibody within 13 days using increasing dosages.

Results

The intraperitoneal treatment with removab® was able to stop the production of ascites in 22 of 23 patients. These patients were ascites-free at the end of the study (day 37). Immunocytochemical quantification of tumor cells in the ascites fluid showed a dramatic reduction of EpCAM⁺ cells (>log 5). In addition, clinically significant improvement of the quality of life was observed. The majority of adverse events was mild to moderate. The most common side effects observed in the study were fever (82.6%), nausea (60.9%), vomiting (56.5%), and abdominal pain (30.4%). The MDT (maximal tolerated dose) was reached at the increasing dosages of 10 µg–20 µg–50 µg–200 µg–200 µg.

Conclusion

In conclusion, intraperitoneal treatment with removab® was safe, well tolerated and showed encouraging efficacy in patients with malignant ascites due to ovarian cancer. Thus, the new concept of the anti-EpCAM x anti-CD3 anti-

body might offer a promising treatment option for patients with epithelial tumors.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

